



Public Data-based Report



Validation Report Service

Cryo-EM Map Validation Report

Report to assess Cryo-EM Volume Map at Level(s) 0, 1

This report has been generated based on data publicly available at [EMDB](#).

Basic Entry Information:

EMDB ID: [EMD-64625](#)

Title: Cryo-EM structure of the 48-nm repeat doublet microtubule from Tssk6 KO mouse sperm

Authors: [See EMDB entry link](#)

Deposited on: 2025-05-16T00:00:00

Reported Resolution: 4.5 Å

Contact Us:

Instruct Image Processing Center ([I²PC](#))
Biocomputing Unit ([BCU](#))
i2pc@cnb.csic.es
[VRS Website](#)

National Center for Biotechnology (CNB)
St/ Darwin, 3 (Autonomous University of Madrid)
28049 Cantoblanco, Madrid (Spain)

Last update: **August 21, 2026, 1:08pm**

Context

Cryo-electron microscopy is currently one of the most active techniques in Structural Biology. The number of maps deposited at the [Electron Microscopy Data Bank](#) is rapidly growing every year and keeping the quality of the submitted maps is essential to maintain the scientific quality of the field. The ultimate quality measure is the consistency of the map and an atomic model. However, this is only possible for high resolution maps. Over the years there have been many suggestions about validation measures of 3DEM maps. Unfortunately, most of these measures are not currently in use for their spread in multiple software tools and the associated difficulty to access them. To alleviate this problem, we made available a validation grading system that evaluate the information provided to assess the map.

This system grades a map from 0 to 5 depending on the amount of information available. In this way, a map could be validated at Level 0 (the deposited map), 1 (two half maps), 2 (2D classes), 3 (particles), 4 (... + angular assignment), 5 (... + micrographs and coordinates). In addition, we can have three optional qualifiers: A (... + atomic model), W (... + image processing workflow), and O (... + other techniques). To know more about this service read this [paper](#)

This Validation Report Service uses Scipion (see this [link](#) for more detail) as workflow engine and ChimeraX (see this [link](#) for more detail) to generate the 3D views. For more information about the different methods and softwares used for this report, see the references [here](#).



Summarized overall quality

The map seems to be well centered. There seems to be a problem with the suggested threshold (see Sec. [2.2](#)). There seems to be a problem with the map's background (see Sec. [2.3](#)).

The overall score (passing tests) of this report is 2 out of 4 evaluable items.

0.a Mass analysis	Sec. 2.1	OK
0.b Mask analysis	Sec. 2.2	2 warnings
0.c Background analysis	Sec. 2.3	2 warnings
0.d B-factor analysis	Sec. 2.4	OK

Summary of the warnings across sections.

Section 2.2 (0.b Mask analysis)

1. **There might be a problem of connectivity at this threshold because more than 5 connected components are needed to reach 95% of the total mask. Probably a smaller threshold will not cause this issue.**
2. **There might be a problem in the construction of the mask, because the overlap is smaller than 0.75. A common reason is that the suggested threshold causes too many disconnected components.**

Section 2.3 (0.c Background analysis)

1. **The null hypothesis that the background mean is 0 has been rejected because the p-value of the comparison is smaller than 0.001**
2. **There is a significant proportion of outlier values in the background (cdf5 ratio=7950.46)**

Contents

1	Input data	6
2	Level 0 analysis	13
2.1	Level 0.a Mass analysis	13
2.2	Level 0.b Mask analysis	14
2.3	Level 0.c Background analysis	19
2.4	Level 0.d B-factor analysis	21
3	Half maps	23

1 Input data

Input map: emd_64625.map

SHA256 hash: 95f73ec909e492c9bbb37fa1a85b1e80198c7b5bbeb3f32bf4bd7c182c3a4a9

Voxel size: 0.930000 (Å)

Visualization threshold: 0.004000

Resolution estimated by user: 4.5


Orthogonal slices of the input map

Explanation:


In the orthogonal slices of the map, the noise outside the protein should not have any structure (stripes going out, small blobs, particularly high or low densities, ...)

Results:


See Fig. 1.



(a) X Slice 382



(b) Y Slice 382




(c) Z Slice 382

Figure 1: Central slices of the input map in the three dimensions


Orthogonal slices of maximum variance of the input map

Results:


See Fig. 2.



(a) X Slice 388



(b) Y Slice 382



(c) Z Slice 382

Figure 2: Slices of maximum variation in the three dimensions


Orthogonal projections of the input map

Explanation:


In the projections there should not be stripes (this is an indication of directional overweighting, or angular attraction), and there should not be a dark halo around or inside the structure (this is an indication of incorrect CTF correction or the reconstruction of a biased map).

Results:


See Fig. [3](#).



(a) X Projection



(b) Y Projection



(c) Z Projection

Figure 3: Projections in the three dimensions

Isosurface views of the input map

Explanation:

An isosurface is the surface of all points that have the same gray value. In these views there should not be many artifacts or noise blobs around the map.

Results:

See Fig. 4.




Figure 4: Isosurface at threshold=0.004000. Views generated by ChimeraX at a the following X, Y, Z angles: View 1 (0, -90, -90), View 2 (-90, 0, -90), View 3 (0, 0, 0).


Orthogonal slices of maximum variance of the mask with hard borders

Explanation:


The mask with hard borders has been calculated at the suggested threshold 0.004000, the largest connected component was selected, and then dilated by 2Å.

Results:


See Fig. 5.



(a) X Slice 390



(b) Y Slice 271



(c) Z Slice 421

Figure 5: Slices of maximum variation in the three dimensions of the mask with hard borders


Orthogonal slices of maximum variance of the mask with soft borders

Explanation:

The mask with soft borders has been calculated at the suggested threshold 0.004000, the largest connected component was selected, and then dilated by

2Å.


Results:
See Fig. 6.



(a) X Slice 388



(b) Y Slice 258



(c) Z Slice 422

Figure 6: Slices of maximum variation in the three dimensions of the mask with soft borders

2 Level 0 analysis

2.1 Level 0.a Mass analysis

Explanation:

The reconstructed map must be relatively well centered in the box, and there should be at least $30\text{\AA}$ (the exact size depends on the CTF) on each side to make sure that the CTF can be appropriately corrected.

Results:

The space from the left and right in X are 132.06 and 164.61 $\text{\AA}$, respectively. There is a decentering ratio $(\text{abs}(\text{Right-Left})/\text{Size})\%$ of 4.58%

The space from the left and right in Y are 202.74 and 236.22 $\text{\AA}$, respectively. There is a decentering ratio $(\text{abs}(\text{Right-Left})/\text{Size})\%$ of 4.71%

The space from the left and right in Z are 51.15 and 53.01 $\text{\AA}$, respectively. There is a decentering ratio $(\text{abs}(\text{Right-Left})/\text{Size})\%$ of 0.26%

The center of mass is at (x,y,z)=(376.04,418.55,383.39). The decentering of the center of mass (abs(Center)/Size)% is 0.78, 4.78, and 0.18, respectively.

Automatic criteria: The validation is OK if 1) the decentering and center of mass less than 20% of the map dimensions in all directions, and 2) the extra space on each direction is more than 20% of the map dimensions. For local reconstruction, focused refinement, or similar, warnings are expected.

STATUS: OK

2.2 Level 0.b Mask analysis

Explanation:

The map at the suggested threshold should have most of its mass concentrated in a single connected component. It is normal that after thresholding there are a few thousands of very small, disconnected noise blobs. However, there total mass should not exceed 10%. The raw mask (just thresholding) and the mask constructed for the analysis (thresholding + largest connected component + dilation) should significantly overlap. Overlap is defined by the overlapping coefficient (size(Raw AND Constructed)/size(Raw)) that is a number between 0 and 1, the closer to 1, the more they agree.

Results:

Raw mask: At threshold 0.004000, there are 35083 connected components with a total number of voxels of 4685936 and a volume of 3769165.42 Å³ (see Fig. 7). The size and percentage of the total number of voxels for the raw mask are listed below (up to 95% of the mass or the first 100 clusters, whatever happens first), the list contains (No. voxels (volume in Å³), percentage, cumulated percentage):

(2223131 (1788190.98), 47.44, 47.44)(110602 (88963.49), 2.36, 49.80)(84526 (67989.08), 1.80, 51.61)(50323 (40477.66), 1.07, 52.68)(32164 (25871.34), 0.69, 53.37)(20149 (16206.99), 0.43, 53.80)(17922 (14415.69), 0.38, 54.18)(17165 (13806.79), 0.37, 54.55)(15212 (12235.88), 0.32, 54.87)(13190 (10609.47),


0.28, 55.15)(12151 (9773.74), 0.26, 55.41)(11234 (9036.15), 0.24, 55.65)(10366 (8337.96), 0.22, 55.87)(10014 (8054.83), 0.21, 56.09)(9906 (7967.96), 0.21, 56.30)(9375 (7540.85), 0.20, 56.50)(8838 (7108.91), 0.19, 56.69)(8733 (7024.45), 0.19, 56.87)(8673 (6976.19), 0.19, 57.06)(8579 (6900.58), 0.18, 57.24)(8236 (6624.68), 0.18, 57.42)(7782 (6259.51), 0.17, 57.58)(7590 (6105.07), 0.16, 57.74)(7541 (6065.66), 0.16, 57.91)(7259 (5838.83), 0.15, 58.06)(7183 (5777.70), 0.15, 58.21)(7129 (5734.26), 0.15, 58.37)(6475 (5208.21), 0.14, 58.50)(6350 (5107.67), 0.14, 58.64)(6319 (5082.73), 0.13, 58.77)(6303 (5069.86), 0.13, 58.91)(6158 (4953.23), 0.13, 59.04)(6115 (4918.64), 0.13, 59.17)(6024 (4845.45), 0.13, 59.30)(5983 (4812.47), 0.13, 59.43)(5963 (4796.38), 0.13, 59.55)(5910 (4753.75), 0.13, 59.68)(5774 (4644.36), 0.12, 59.80)(5662 (4554.27), 0.12, 59.92)(5565 (4476.25), 0.12, 60.04)(5452 (4385.35), 0.12, 60.16)(5292 (4256.66), 0.11, 60.27)(5267 (4236.55), 0.11, 60.38)(5182 (4168.18), 0.11, 60.50)(5102 (4103.83), 0.11, 60.60)(5076 (4082.92), 0.11, 60.71)(4522 (3637.30), 0.10, 60.81)(4479 (3602.72), 0.10, 60.90)(4312 (3468.39), 0.09, 61.00)(4126 (3318.78), 0.09, 61.08)(4107 (3303.49), 0.09, 61.17)(4079 (3280.97), 0.09, 61.26)(4056 (3262.47), 0.09, 61.35)(4009 (3224.67), 0.09, 61.43)(3994 (3212.60), 0.09, 61.52)(3813 (3067.01), 0.08, 61.60)(3810 (3064.60), 0.08, 61.68)(3789 (3047.71), 0.08, 61.76)(3702 (2977.73), 0.08, 61.84)(3697 (2973.71), 0.08, 61.92)(3692 (2969.69), 0.08, 62.00)(3630 (2919.82), 0.08, 62.07)(3608 (2902.12), 0.08, 62.15)(3597 (2893.27), 0.08, 62.23)(3549 (2854.66), 0.08, 62.30)(3547 (2853.05), 0.08, 62.38)(3451 (2775.84), 0.07, 62.45)(3407 (2740.44), 0.07, 62.53)(3398 (2733.21), 0.07, 62.60)(3306 (2659.20), 0.07, 62.67)(3281 (2639.10), 0.07, 62.74)(3276 (2635.07), 0.07, 62.81)(3270 (2630.25), 0.07, 62.88)(3179 (2557.05), 0.07, 62.95)(3103 (2495.92), 0.07, 63.01)(3054 (2456.51), 0.07, 63.08)(3008 (2419.51), 0.06, 63.14)(2974 (2392.16), 0.06, 63.21)(2962 (2382.51), 0.06, 63.27)(2948 (2371.24), 0.06, 63.33)(2947 (2370.44), 0.06, 63.39)(2921 (2349.53), 0.06, 63.46)(2897 (2330.22), 0.06, 63.52)(2891 (2325.40), 0.06, 63.58)(2871 (2309.31), 0.06, 63.64)(2846 (2289.20), 0.06, 63.70)(2841 (2285.18), 0.06, 63.76)(2801 (2253.00), 0.06, 63.82)(2781 (2236.92), 0.06, 63.88)(2688 (2162.11), 0.06, 63.94)(2664 (2142.81), 0.06, 64.00)(2588 (2081.68), 0.06, 64.05)(2579 (2074.44), 0.06, 64.11)(2569 (2066.39), 0.05, 64.16)(2568 (2065.59), 0.05, 64.22)(2536 (2039.85), 0.05, 64.27)(2527 (2032.61), 0.05, 64.32)(2505 (2014.91), 0.05, 64.38)(2488 (2001.24), 0.05, 64.43)

Number of components to reach 95% of the mass: 7229


The average size of the remaining 27854 components is 8.41 voxels (0.80

$\text{\AA}^3$). Their size go from 38 voxels ($30.57 \text{ \AA}^3$) to 1 voxels ($0.80 \text{ \AA}^3$).


The slices of the raw mask can be seen in Fig. 7.



(a) X Slice 384



(b) Y Slice 271



(c) Z Slice 382

Figure 7: Maximum variance slices in the three dimensions of the raw mask

The following table shows the variation of the mass enclosed at different thresholds (see Fig. 8):

Threshold	Voxel mass	Molecular mass(kDa)	# Aminoacids
0.0005	54580489.00	36372.99	330663.54
0.0010	35884491.00	23913.79	217398.07
0.0015	24234503.00	16150.12	146819.25
0.0021	17124242.00	11411.77	103743.35
0.0026	12394069.00	8259.53	75086.66
0.0031	8927895.00	5949.64	54087.63
0.0036	6285371.00	4188.63	38078.50
0.0041	4297796.00	2864.09	26037.22
0.0046	2848353.00	1898.17	17256.10
0.0051	1827836.00	1218.09	11073.53
0.0057	1135134.00	756.46	6876.95
0.0062	681018.00	453.84	4125.79
0.0067	393327.00	262.12	2382.88
0.0072	217569.00	144.99	1318.09
0.0077	114702.00	76.44	694.90
0.0082	57485.00	38.31	348.26
0.0087	26955.00	17.96	163.30
0.0093	11874.00	7.91	71.94
0.0098	4841.00	3.23	29.33
0.0103	1843.00	1.23	11.17
0.0108	667.00	0.44	4.04
0.0113	233.00	0.16	1.41
0.0118	62.00	0.04	0.38
0.0123	14.00	0.01	0.08
0.0129	0.00	0.00	0.00




Figure 8: Voxel mass as a function of the gray level.

Constructed mask: After keeping the largest component of the previous mask and dilating it by $2\text{\AA}$, there is a total number of voxels of 15859422 and a volume of $12756637.10 \text{ \AA}^3$. The overlap between the raw and constructed mask is 0.48.

Automatic criteria: The validation is OK if 1) to keep 95% of the mass we need to keep at most 5 connected components; and 2) the average volume of the blobs outside the given threshold has a size smaller than $5\text{\AA}^3$; and 3) the overlap between the raw mask and the mask constructed for the analysis is larger than 75%.

WARNINGS: 2 warnings

1. **There might be a problem of connectivity at this threshold because more than 5 connected components are needed to reach 95% of the total mask. Probably a smaller threshold will not cause this issue.**
2. **There might be a problem in the construction of the mask, because the overlap is smaller than 0.75. A common reason is that the suggested threshold causes too many disconnected**

components.

2.3 Level 0.c Background analysis

Explanation:


Background is defined as the region outside the macromolecule mask. The background mean should be zero, and the number of voxels with a very low or very high value (below 5 standard deviations of the noise) should be very small and they should be randomly distributed without any specific structure. Sometimes, you can see some structure due to the symmetry of the structure.

Results:


The null hypothesis that the background mean is 0 was tested with a one-sample Student's t-test. The resulting t-statistic and p-value were -512.84 and 0.000000, respectively.

The mean and standard deviation (sigma) of the background were -0.000021 and 0.000851. The percentage of background voxels whose absolute value is larger than 5 times the standard deviation is 0.46 % (see Fig. 9). The same percentage from a Gaussian would be 0.000057% (ratio between the two percentages: 7950.461874).


Slices of the background beyond 5*sigma can be seen in Fig. 9.



(a) X Slice 384



(b) Y Slice 513



(c) Z Slice 604

Figure 9: Maximum variance slices in the three dimensions of the parts of the background beyond 5σ

Automatic criteria: The validation is OK if 1) the p-value of the null hypothesis that the background has 0 mean is larger than 0.001; and 2) the number of voxels above or below 5 sigma is smaller than 20 times the amount

expected for a Gaussian with the same standard deviation whose mean is 0.

WARNINGS: 2 warnings

1. **The null hypothesis that the background mean is 0 has been rejected because the p-value of the comparison is smaller than 0.001**
2. **There is a significant proportion of outlier values in the background (cdf5 ratio=7950.46)**

2.4 Level 0.d B-factor analysis

Explanation:

The B-factor line (see this [link](#) for more details) fitted between 15Å and the resolution reported should have a slope that is between 0 and 300 Å².

Results:

Fig. 10 shows the logarithm (in natural units) of the structure factor (the module squared of the Fourier transform) of the experimental map, its fitted line, and the corrected map. The estimated B-factor was -68.3. The fitted line was $\log(|F|^2) = -17.1/R^2 + (-15.4)$.







Figure 10: Guinier plot. The X-axis is the square of the inverse of the resolution in $\text{\AA}$.





(c) Z Slice 382

Figure 11: Slices of maximum variation in the three dimensions of the B-factor corrected map

Automatic criteria: The validation is OK if the B-factor is in the range $[-300,0]$.

STATUS: OK

3 Half maps

Half map 1: emd_64625_half_map_1.map

SHA256 hash: 7f35010a9475f5363cdf0162fab1096d42a18ebd2f8057cde7bfac6dd11162ef


Half map 2: emd_64625_half_map_2.map

SHA256 hash: 721713d64febe5ee43362a1b5412b3d21c0b9cf689bcbfc5fe74647c4b7c8a11


Slices of the first half map can be seen in Fig. 12.

Slices of the second half map can be seen in Fig. 13.


Slices of the difference between both maps can be seen in Fig. 14. There should not be any structure in this difference. Sometimes some patterns are seen if the map is symmetric.



(a) X Slice 386




(b) Y Slice 381




(c) Z Slice 390


Figure 12: Slices of maximum variation in the three dimensions of Half 1



(a) X Slice 393




(b) Y Slice 381




(c) Z Slice 393


Figure 13: Slices of maximum variation in the three dimensions of Half 2



(a) X Slice 378



(b) Y Slice 396



(c) Z Slice 389

Figure 14: Slices of maximum variation in the three dimensions of the difference Half1-Half2.