



Public Data-based Report



Validation Report Service

Cryo-EM Map Validation Report

Report to assess Cryo-EM Volume Map at Level(s) 1, 0

This report has been generated based on data publicly available at [EMDB](#).

Basic Entry Information:

EMDB ID: [EMD-71628](#)

Title: Human SRCAP-nucleosome complex in the fully-engaged state, with unbound GAS41 (unmasked refinement, filtered by local resolution)

Authors: [See EMDB entry link](#)

Deposited on: 2025-07-07T00:00:00

Reported Resolution: 6.8 Å

Contact Us:

Instruct Image Processing Center ([I²PC](#))
Biocomputing Unit ([BCU](#))
i2pc@cnb.csic.es
[VRS Website](#)

National Center for Biotechnology (CNB)
St/ Darwin, 3 (Autonomous University of Madrid)
28049 Cantoblanco, Madrid (Spain)

Last update: **September 4, 2026, 10:39pm**

Context

Cryo-electron microscopy is currently one of the most active techniques in Structural Biology. The number of maps deposited at the [Electron Microscopy Data Bank](#) is rapidly growing every year and keeping the quality of the submitted maps is essential to maintain the scientific quality of the field. The ultimate quality measure is the consistency of the map and an atomic model. However, this is only possible for high resolution maps. Over the years there have been many suggestions about validation measures of 3DEM maps. Unfortunately, most of these measures are not currently in use for their spread in multiple software tools and the associated difficulty to access them. To alleviate this problem, we made available a validation grading system that evaluate the information provided to assess the map.

This system grades a map from 0 to 5 depending on the amount of information available. In this way, a map could be validated at Level 0 (the deposited map), 1 (two half maps), 2 (2D classes), 3 (particles), 4 (... + angular assignment), 5 (... + micrographs and coordinates). In addition, we can have three optional qualifiers: A (... + atomic model), W (... + image processing workflow), and O (... + other techniques). To know more about this service read this [paper](#)

This Validation Report Service uses Scipion (see this [link](#) for more detail) as workflow engine and ChimeraX (see this [link](#) for more detail) to generate the 3D views. For more information about the different methods and softwares used for this report, see the references [here](#).



Summarized overall quality

The map seems to have some problem in its centering or extra space (see Sec. [2.1](#)). There is no problem with the suggested threshold. There seems to be a problem with the map's background (see Sec. [2.3](#)). There seems to be a problem with its B-factor (see Sec. [2.4](#)).

The overall score (passing tests) of this report is 1 out of 4 evaluable items.

0.a Mass analysis	Sec. 2.1	2 warnings
0.b Mask analysis	Sec. 2.2	OK
0.c Background analysis	Sec. 2.3	2 warnings
0.d B-factor analysis	Sec. 2.4	1 warnings

Summary of the warnings across sections.

Section 2.1 (0.a Mass analysis)

1. **The center of mass in Y may be significantly shifted. This is common when the refinement is applied exclusively to one protein region.**
2. **The center of mass in Z may be significantly shifted. This is common when the refinement is applied exclusively to one protein region.**

Section 2.3 (0.c Background analysis)

1. **The null hypothesis that the background mean is 0 has been rejected because the p-value of the comparison is smaller than 0.001**
2. **There is a significant proportion of outlier values in the background (cdf5 ratio=16920.28)**

Section 2.4 (0.d B-factor analysis)

1. **The B-factor is out of the interval [-300,0].**

Contents

1	Input data	6
2	Level 0 analysis	13
2.1	Level 0.a Mass analysis	13
2.2	Level 0.b Mask analysis	14
2.3	Level 0.c Background analysis	18
2.4	Level 0.d B-factor analysis	20
3	Half maps	23

1 Input data

Input map: emd_71628.map

SHA256 hash: 80026da4e2118c17d7e61e84ce38b10dd4f1624eed7ee8a6fe2c47a249600bf7

Voxel size: 1.025000 (Å)

Visualization threshold: 0.006500

Resolution estimated by user: 6.8


Orthogonal slices of the input map

Explanation:


In the orthogonal slices of the map, the noise outside the protein should not have any structure (stripes going out, small blobs, particularly high or low densities, ...)

Results:


See Fig. 1.



(a) X Slice 192



(b) Y Slice 192




(c) Z Slice 192

Figure 1: Central slices of the input map in the three dimensions


Orthogonal slices of maximum variance of the input map

Results:


See Fig. 2.



(a) X Slice 191



(b) Y Slice 229



(c) Z Slice 186

Figure 2: Slices of maximum variation in the three dimensions


Orthogonal projections of the input map

Explanation:


In the projections there should not be stripes (this is an indication of directional overweighting, or angular attraction), and there should not be a dark halo around or inside the structure (this is an indication of incorrect CTF correction or the reconstruction of a biased map).

Results:


See Fig. [3](#).



(a) X Projection



(b) Y Projection



(c) Z Projection

Figure 3: Projections in the three dimensions

Isosurface views of the input map

Explanation:

An isosurface is the surface of all points that have the same gray value. In these views there should not be many artifacts or noise blobs around the map.

Results:

See Fig. 4.




Figure 4: Isosurface at threshold=0.006500. Views generated by ChimeraX at a the following X, Y, Z angles: View 1 (0, -90, -90), View 2 (-90, 0, -90), View 3 (0, 0, 0).


Orthogonal slices of maximum variance of the mask with hard borders

Explanation:


The mask with hard borders has been calculated at the suggested threshold 0.006500, the largest connected component was selected, and then dilated by 2Å.

Results:


See Fig. 5.



(a) X Slice 187



(b) Y Slice 203



(c) Z Slice 187

Figure 5: Slices of maximum variation in the three dimensions of the mask with hard borders

Orthogonal slices of maximum variance of the mask with soft borders

Explanation:

The mask with soft borders has been calculated at the suggested threshold 0.006500, the largest connected component was selected, and then dilated by

2Å.

Results:
See Fig. 6.



(a) X Slice 187



(b) Y Slice 203



(c) Z Slice 194

Figure 6: Slices of maximum variation in the three dimensions of the mask with soft borders

2 Level 0 analysis

2.1 Level 0.a Mass analysis

Explanation:

The reconstructed map must be relatively well centered in the box, and there should be at least $30\text{\AA}$ (the exact size depends on the CTF) on each side to make sure that the CTF can be appropriately corrected.

Results:

The space from the left and right in X are 106.60 and 72.77 $\text{\AA}$, respectively. There is a decentering ratio $(\text{abs}(\text{Right-Left})/\text{Size})\%$ of 8.59%

The space from the left and right in Y are 97.37 and 103.52 $\text{\AA}$, respectively. There is a decentering ratio $(\text{abs}(\text{Right-Left})/\text{Size})\%$ of 1.56%

The space from the left and right in Z are 86.10 and 114.80 $\text{\AA}$, respectively. There is a decentering ratio $(\text{abs}(\text{Right-Left})/\text{Size})\%$ of 7.29%

The center of mass is at (x,y,z)=(155.06,-253.55,275.99). The decentering of the center of mass (abs(Center)/Size)% is 9.62, 116.03, and 21.87, respectively.

Automatic criteria: The validation is OK if 1) the decentering and center of mass less than 20% of the map dimensions in all directions, and 2) the extra space on each direction is more than 20% of the map dimensions. For local reconstruction, focused refinement, or similar, warnings are expected.

WARNINGS: 2 warnings

1. **The center of mass in Y may be significantly shifted. This is common when the refinement is applied exclusively to one protein region.**
2. **The center of mass in Z may be significantly shifted. This is common when the refinement is applied exclusively to one protein region.**

2.2 Level 0.b Mask analysis

Explanation:

The map at the suggested threshold should have most of its mass concentrated in a single connected component. It is normal that after thresholding there are a few thousands of very small, disconnected noise blobs. However, their total mass should not exceed 10%. The raw mask (just thresholding) and the mask constructed for the analysis (thresholding + largest connected component + dilation) should significantly overlap. Overlap is defined by the overlapping coefficient (size(Raw AND Constructed)/size(Raw)) that is a number between 0 and 1, the closer to 1, the more they agree.

Results:

Raw mask: At threshold 0.006500, there are 12 connected components with a total number of voxels of 1115359 and a volume of 1201119.65 Å³ (see Fig. 7). The size and percentage of the total number of voxels for the raw mask are listed below (up to 95% of the mass or the first 100 clusters, whatever

happens first), the list contains (No. voxels (volume in $\text{\AA}^3$), percentage, cumulated percentage):

(1115246 (1200997.96), 99.99, 99.99)


Number of components to reach 95% of the mass: 1

The average size of the remaining 11 components is 10.27 voxels ($1.08 \text{ \AA}^3$).
Their size go from 1115246 voxels ($1200997.96 \text{ \AA}^3$) to 1 voxels ($1.08 \text{ \AA}^3$).


The slices of the raw mask can be seen in Fig. 7.



(a) X Slice 189



(b) Y Slice 201



(c) Z Slice 194

Figure 7: Maximum variance slices in the three dimensions of the raw mask

The following table shows the variation of the mass enclosed at different thresholds (see Fig. 8):

Threshold	Voxel mass	Molecular mass(kDa)	# Aminoacids
0.0022	1577308.00	1407.28	12793.46
0.0044	1305038.00	1164.36	10585.10
0.0067	1102142.00	983.34	8939.42
0.0089	927759.00	827.75	7525.01
0.0111	769049.00	686.15	6237.72
0.0133	632032.00	563.90	5126.38
0.0155	502433.00	448.27	4075.21
0.0178	388309.00	346.45	3149.55
0.0200	296957.00	264.95	2408.60
0.0222	223289.00	199.22	1811.09
0.0244	162615.00	145.09	1318.96
0.0266	113473.00	101.24	920.37
0.0289	76549.00	68.30	620.88
0.0311	50215.00	44.80	407.29
0.0333	32013.00	28.56	259.66
0.0355	19588.00	17.48	158.88
0.0377	11495.00	10.26	93.24
0.0400	6432.00	5.74	52.17
0.0422	3408.00	3.04	27.64
0.0444	1589.00	1.42	12.89
0.0466	654.00	0.58	5.30
0.0488	225.00	0.20	1.82
0.0511	82.00	0.07	0.67
0.0533	19.00	0.02	0.15




Figure 8: Voxel mass as a function of the gray level.

Constructed mask: After keeping the largest component of the previous mask and dilating it by $2\text{\AA}$, there is a total number of voxels of 1651619 and a volume of $1778613.02\text{ \AA}^3$. The overlap between the raw and constructed mask is 1.00.

Automatic criteria: The validation is OK if 1) to keep 95% of the mass we need to keep at most 5 connected components; and 2) the average volume of the blobs outside the given threshold has a size smaller than $5\text{\AA}^3$; and 3) the overlap between the raw mask and the mask constructed for the analysis is larger than 75%.

STATUS: OK

2.3 Level 0.c Background analysis

Explanation:

Background is defined as the region outside the macromolecule mask. The background mean should be zero, and the number of voxels with a very low or very high value (below 5 standard deviations of the noise) should be very


small and they should be randomly distributed without any specific structure. Sometimes, you can see some structure due to the symmetry of the structure.

Results:


The null hypothesis that the background mean is 0 was tested with a one-sample Student's t-test. The resulting t-statistic and p-value were -2864.90 and 0.000000, respectively.

The mean and standard deviation (σ) of the background were -0.000363 and 0.000939. The percentage of background voxels whose absolute value is larger than 5 times the standard deviation is 0.97 % (see Fig. 9). The same percentage from a Gaussian would be 0.000057% (ratio between the two percentages: 16920.275998).


Slices of the background beyond 5σ can be seen in Fig. 9.



(a) X Slice 149



(b) Y Slice 189



(c) Z Slice 205

Figure 9: Maximum variance slices in the three dimensions of the parts of the background beyond 5*sigma

Automatic criteria: The validation is OK if 1) the p-value of the null hypothesis that the background has 0 mean is larger than 0.001; and 2) the number of voxels above or below 5 sigma is smaller than 20 times the amount expected for a Gaussian with the same standard deviation whose mean is 0.

WARNINGS: 2 warnings

1. **The null hypothesis that the background mean is 0 has been rejected because the p-value of the comparison is smaller than 0.001**
2. **There is a significant proportion of outlier values in the background (cdf5 ratio=16920.28)**

2.4 Level 0.d B-factor analysis

Explanation:

The B-factor line (see this [link](#) for more details) fitted between 15Å and the resolution reported should have a slope that is between 0 and 300 Å².

Results:

Fig. 10 shows the logarithm (in natural units) of the structure factor (the module squared of the Fourier transform) of the experimental map, its fitted line, and the corrected map. The estimated B-factor was -439.2. The fitted line was $\log(|F|^2) = -109.8/R^2 + (-12.5)$.




Figure 10: Guinier plot. The X-axis is the square of the inverse of the resolution in Å.




Figure 11: Slices of maximum variation in the three dimensions of the B-factor corrected map

Automatic criteria: The validation is OK if the B-factor is in the range $[-300,0]$.

WARNINGS: 1 warnings

1. **The B-factor is out of the interval [-300,0].**

3 Half maps

Half map 1: emd_71628_half_map_1.map

SHA256 hash: 33fd14d5fb2077f56e5b8a87d1f5d81dcbe42f65bccee0f2e34fc4f42f6f2b27


Half map 2: emd_71628_half_map_2.map

SHA256 hash: 9f9d7540ffa18fde5425ac3e52151edc8bddd4dafab0fffe80bb271cec549b


Slices of the first half map can be seen in Fig. 12.

Slices of the second half map can be seen in Fig. 13.


Slices of the difference between both maps can be seen in Fig. 14. There should not be any structure in this difference. Sometimes some patterns are seen if the map is symmetric.



(a) X Slice 193




(b) Y Slice 202




(c) Z Slice 196


Figure 12: Slices of maximum variation in the three dimensions of Half 1



(a) X Slice 189




(b) Y Slice 195




(c) Z Slice 186


Figure 13: Slices of maximum variation in the three dimensions of Half 2



(a) X Slice 184



(b) Y Slice 186



(c) Z Slice 200

Figure 14: Slices of maximum variation in the three dimensions of the difference Half1-Half2.